




CMV-specific immune monitoring test with QuantiFERON[®]-CMV

"How do I know which of my transplant patients are at risk of CMV disease?"



Which transplant patients are at risk of CMV disease post-prophylaxis?

Current immunosuppressive therapies used to prevent the rejection of a transplanted organ have detrimental effects upon the T-lymphocytes and cell-mediated immune responses in solid organ transplant (SOT) recipients. These patients have an increased susceptibility to viral infections post-transplant.^{2, 3} Approximately half of these patients show signs of active Cytomegalovirus (CMV) infection (ie. viral replication) after completion of antiviral prophylaxis.¹

Consequently, CMV-associated morbidity and mortality in transplant recipients are very high.³⁻⁵ Infection can result not only in organ-specific conditions affecting the brain, lungs, and liver, but also in opportunistic infections, increased allograft rejection, and patient death. These CMV-related complications add an estimated 49% to the cost of transplant care.⁶

The immune status of the transplant recipient can influence the (re)activation of CMV in transplant recipients. A specific cytokine marker for cellular immune responses, interferon-gamma (IFN- γ) plays a key role.^{7, 8} Secreted from CMV-specific T-cells in response to antigens associated with CMV infection,^{8, 9} IFN- γ levels may indicate a patient's overall level of cell-mediated immunity.

Cytomegalovirus (CMV) is the most common and problematic viral infection in solid organ transplant recipients.

How can QuantiFERON-CMV help identify at-risk patients?

QuantiFERON-CMV uses specialized (1 mL) blood collection tubes that are coated with peptides simulating CD8⁺-specific epitopes of CMV proteins, along with negative and positive control tubes. Stimulation of CD8⁺ T-cells in whole blood with the CMV peptides results in the production of IFN- γ in infected individuals. An enzyme-linked immunosorbent assay (ELISA) is then used to measure the amount of IFN- γ present in plasma from each of the three tubes (Nil control, CMV-antigen, and Mitogen control). A robust IFN- γ response in the CMV antigen tube is indicative of immunity to CMV.¹¹

Immunity to CMV during Post-Transplantation prophylaxis





Figure 1. A theoretical model of QuantiFERON-CMV responses in a post-transplant setting during and/or following prophylaxis and immune suppression.

Freedom from CMV-related events

Patients who have a cellular immune response to CMV at the end of prophylaxis have a significantly lower risk of developing CMV disease than those who do not have a detectable immune response. This indicates that QuantiFERON-CMV may predict the development of late-onset CMV disease in transplant recipients.^{8, 10-13}

Figure 2. Time to development of CMV disease in patients with a QuantiFERON-CMV Reactive result versus a QuantiFERON-CMV Non-Reactive result at the end of prophylaxis. Data reproduced from Kumar *et al.*¹¹



Patients with a positive QuantiFERON-CMV test remain free from CMV disease significantly more often and for longer than patients with a negative QuantiFERON-CMV after cessation of antiviral prophylaxis.

Clinical confidence

QuantiFERON-CMV may assist¹³ your ability to:

- Predict the risk of new and recurrent CMV disease
- Guide therapeutic decision-making
- Improve patient health.

QuantiFERON-CMV is not a direct test for determining CMV infection and should not solely be used to exclude CMV infection.

The accuracy, efficacy, and utility of QuantiFERON-CMV for monitoring CMV-related changes in cell-mediated immunity has been demonstrated in numerous studies.^{8, 10-13}

New international consensus guidelines on the management of CMV in SOT

The recently-published "International Consensus Guidelines on the Management of Cytomegalovirus in solid organ transplantation"¹⁴ suggest that an ideal immune monitoring assay should:

- Assess the quantity and function of a transplant recipient's CD4⁺ and CD8⁺ T-cells
- Be able to measure interferon- γ
- Be simple to perform, cost-effective, and reproducible
- Have a rapid turnaround time
- Allow for specimens to be easily shipped to specialized referral laboratories.

QuantiFERON-CMV meets virtually all the criteria specified by the guidelines above and is the only standardized, commercially-available immune monitoring assay that is specific for CMV.¹⁴

Studies now highlight that monitoring a patient's level of cellular immunity to CMV using QuantiFERON-CMV could help guide the optimal duration of costly anti-CMV prophylaxis in high-risk patients.^{8, 10, 11, 13-16}

About QuantiFERON® Technology




Figure 3. Diagram illustrating the QuantiFERON-CMV process.

QuantiFERON Technology is a unique approach to disease detection and monitoring—the only *in vitro* diagnostic technology available for detection of cell-mediated immune responses from whole blood samples.

Individuals exposed to CMV and other diseases have specific T-cell lymphocytes in their blood. These T-cells are a memory bank of an individual's immune system, recognizing antigens to which the T-cells have been previously exposed. When a disease-specific antigen is combined with the blood of an individual who has been exposed to that disease, a rapid re-stimulation of the T-cells with specific memory of that antigen occurs. These antigen-specific T-cells respond by secreting IFN- γ , which can be measured as a specific marker of immune response against that disease antigen.

QuantiFERON-CMV is not US FDA-approved and is limited to research use only in the United States.

QuantiFERON-CMV is CE Marked for commercial use in Europe.

References

1. Paya C, et al. *Am J Transpl.* 2004; 4:611–20.
2. Smyth RL, et al. *Transplantation.* 1991; 52:480–2.
3. Boeckh M, Nichols WG. *Herpes.* 2003; 10:12–16.
4. Gandhi MK, Khanna R. *Lancet Infect Dis.* 2004; 4:725–38.
5. Humar A, et al. *Transplantation.* 2000; 70:310–3.
6. Kim WR, et al. *Transplantation.* 2000; 69:357–61.
7. Sylwester AW, et al. *J Exp Med.* 2005; 202:673–85.
8. Walker S, et al. *Transpl Infect Dis.* 2007; 9:165–70.
9. Westall G, et al. *Am J Transpl.* 2006; 6:577–84.
10. Westall GP, et al. *Am J Transpl.* 2008; 8:1749–54.
11. Kumar D, et al. *Am J Transpl.* 2009; 9:1214–22.
12. Singh N. *J Clin Virol.* 2006; 35:474–7.
13. Danziger-Isakov LA, et al. *Transplantation.* 2003; 75:1538–43.
14. Kotton CN, et al. *Transplantation* 2010; 89.
15. Danziger-Isakov L, Heeger PS. *Am J Transpl.* 2009; 9:1214–22.
16. Crough T, Khanna R. *Clin Micro Reviews.* 2009; 22:76–98.

© QuantiFERON © QIAGEN Feb 2013 1095925 QM01925003E



Legal Manufacturer: QIAGEN
19300 Germantown Road Germantown,
MD 20874
USA
Tollfree: 1-800-426-8157

Ordering www.qiagen.com/contact | Technical Support support.qiagen.com | Website www.qiagen.com